

References

- [1] E. Kiritsis, “*String theory in a nutshell*,” *Princeton, USA: Univ. Pr. (2007) 588 p*, chapter 13.
- [2] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “*Large N field theories, string theory and gravity*,” *Phys. Rept.* **323** (2000) 183 [ArXiv:hep-th/9905111].
- [3] E. Witten, “*Anti-de Sitter space, thermal phase transition, and confinement in gauge theories*,” *Adv. Theor. Math. Phys.* **2** (1998) 505 [ArXiv:hep-th/9803131];
- [4] T. Sakai and S. Sugimoto, “*Low energy hadron physics in holographic QCD*,” *Prog. Theor. Phys.* **113** (2005) 843 [ArXiv:hep-th/0412141];
“*More on a holographic dual of QCD*,” *Prog. Theor. Phys.* **114** (2006) 1083 [ArXiv:hep-th/0507073].
- [5] O. Aharony, J. Sonnenschein and S. Yankielowicz, “*A holographic model of deconfinement and chiral symmetry restoration*,” *Annals Phys.* **322** (2007) 1420 [ArXiv:hep-th/0604161].
- [6] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, “*QCD and a holographic model of hadrons*,” *Phys. Rev. Lett.* **95**, 261602 (2005) [ArXiv:hep-ph/0501128].
- [7] L. Da Rold and A. Pomarol, “*Chiral symmetry breaking from five dimensional spaces*,” *Nucl. Phys. B* **721**, 79 (2005) [ArXiv:hep-ph/0501218].

- [8] R. Casero, E. Kiritsis and A. Paredes, “*Chiral symmetry breaking as open string tachyon condensation,*” Nucl. Phys. B **787** (2007) 98; [ArXiv:hep-th/0702155].
- [9] U. Gursoy and E. Kiritsis, “Exploring improved holographic theories for QCD: Part I,” JHEP **0802** (2008) 032 [ArXiv:0707.1324][hep-th].
- [10] U. Gursoy, E. Kiritsis and F. Nitti, “Exploring improved holographic theories for QCD: Part II,” JHEP **0802** (2008) 019 [ArXiv:0707.1349][hep-th].
- [11] U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, “*Deconfinement and Gluon Plasma Dynamics in Improved Holographic QCD,*” [ArXiv:0804.0899][hep-th].