The influence of the LMC on dark matter direct detection

Nassim Bozorgnia



IRCHEP 1402, IPM, Iran 21 November 2023



Canada Research Chaires de recherche Chairs du Canada



The Large Magellanic Cloud

The LMC is the most massive satellite of the Milky Way and likely on its first passage around the Galaxy.





Gaia's EDR3 sky map. Credit: ESA/Gaia/DPAC

The effect of the LMC

The LMC introduces perturbations in the DM and stellar halo.



 \mathbf{a}



Conroy et al, Nature 592, 534–536 (2021)

Stellar halo

Local dark matter distribution

Signals in direct DM searches strongly depend on the DM distribution in the Solar neighborhood.



Direct detection event rate

• The differential event rate (per unit detector mass):

$$\frac{dR}{dE_R} = \frac{\rho_{\chi}}{m_{\chi}m_N} \int_{v > v_{\min}} d^3v \ \frac{d\sigma_{\chi N}}{dE_R} \ v \ f_{det}(\mathbf{v}, t)$$

 $v_{\min} = \sqrt{m_N E_R / (2\mu_{\chi N}^2)}$: minimum DM speed required to produce a recoil energy E_R .

Direct detection event rate

• The differential event rate (per unit detector mass):



 $v_{\min} = \sqrt{m_N E_R / (2\mu_{\chi N}^2)}$: minimum DM speed required to produce a recoil energy E_R .

- Astrophysical inputs:
 - **local DM density:** *normalization in event rate.*
 - **local DM velocity distribution:** enters the event rate through an integration.

Direct detection status



Assumption for the DM distribution: Standard Halo Model

 The simplest model for the DM distribution in our Galaxy is the Standard Halo Model (SHM): isothermal sphere with an isotropic Maxwell-Boltzmann velocity distribution.

Drukier, Freese, Spergel, 1986







- Most probable DM speed: $v_c = 220 \text{ km/s}$
- Local DM velocity distribution:

$$f_{\text{gal}}(\mathbf{v}) = \begin{cases} N \exp\left(-\mathbf{v}^2/v_c^2\right) & v < v_{\text{esc}} \\ 0 & v \ge v_{\text{esc}} \end{cases}$$



- Most probable DM speed: $v_c = 220 \text{ km/s}$
- Local DM velocity distribution:

$$f_{\text{gal}}(\mathbf{v}) = \begin{cases} N \exp\left(-\frac{\mathbf{v}^2}{v_c^2}\right) & v < v_{\text{esc}} \\ 0 & v \ge v_{\text{esc}} \end{cases}$$

The actual DM distribution may deviate substantially from the SHM.

The local DM halo has both smooth and un-virialized components:

The local DM halo has both smooth and un-virialized components:



Does the smooth component have a Maxwellian velocity distribution?

The local DM halo has both smooth and un-virialized components:



Extract the DM distribution from cosmological simulations:



Extract the DM distribution from cosmological simulations:



Maxwellian distribution provides a good fit to the DM velocity distribution of Milky Way-like halos in cosmological simulations.

Extract the DM distribution from cosmological simulations:



Rahimi, Vienneau, Bozorgnia, Robertson, 2023 (SIDM EAGLE)

Maxwellian distribution provides a good fit to the DM velocity distribution of Milky Way-like halos in cosmological simulations.

Extract the DM distribution from cosmological simulations:



Effect of LMC on direct detection

The LMC could perturb the high speed tail of the local DM velocity distribution. → Affects direct detection implications for low mass DM.

Effect of LMC on direct detection

- The LMC could perturb the high speed tail of the local DM velocity distribution.
 Affects direct detection implications for low mass DM.

 Besla et al, JCAP 11, 013 (2019)
 Donaldson et al, MNRAS 513, 1, 46 (2022)
- Studied in specially designed idealized simulations.



Besla et al, JCAP 11, 013 (2019)

Effect of LMC on direct detection

- The LMC could perturb the high speed tail of the local DM velocity distribution.
 Affects direct detection implications for low mass DM.

 Besla et al, JCAP 11, 013 (2019)
 Donaldson et al, MNRAS 513, 1, 46 (2022)
- Studied in specially designed idealized simulations.

Are these findings valid for fully cosmological halos with multiple accretion events over their formation history?



Besla et al, JCAP 11, 013 (2019)

- State-of-the-art cosmological magnetohydrodynamical zoom-in simulations of Milky Way size halos.
- 30 halos at the standard resolution:

$m_{\rm DM}~[{ m M}_\odot]$	$m_{ m b}~[{ m M}_{\odot}]$	e [pc]
3×10^{5}	5×10^{4}	369



 Identify I5 Milky Way-LMC analogues based on LMC's stellar mass and distance from host at first pericenter approach.



- Identify I5 Milky Way-LMC analogues based on LMC's stellar mass and distance from host at first pericenter approach.
- Focus on one halo and study the impact of the LMC on the local DM distribution at different times (snapshots) in its orbit.



- Identify I5 Milky Way-LMC analogues based on LMC's stellar mass and distance from host at first pericenter approach.
- Focus on one halo and study the impact of the LMC on the local DM distribution at different times (snapshots) in its orbit.



• Consider four representative snapshots:

Snapshot	Snapshot Description		r _{LMC} [kpc]
lso.	Isolated MW analogue	-2.83	384
Peri.	LMC's 1st pericenter approach	-0.133	32.9
Present day MW-LMC analogu		0	50.6
Fut.	Future MW-LMC analogue	0.175	80.3

Matching the Sun-LMC geometry

 The LMC is predominately moving in the opposite direction of the Solar motion.
 Large relative speeds of DM particles originating from the LMC with respect to the sun.



 Choose the position of the Sun in the simulations such that it matches the observed Sun-LMC geometry.

Local dark matter density

19. 19.	Halo ID	$M_{\mathrm{Infall}}^{\mathrm{LMC}} \ [10^{11} \mathrm{~M_{\odot}}]$	$\rho_{\chi} \ [\text{GeV/cm}^3]$	$\kappa_{\rm LMC}$ [%]	
	1	0.31	0.21	0.14	Percentage of DM particles in
	2	0.31	0.23	0.64	the Solar region
	3	0.34	0.35	0.026	originating from
	4	0.82	0.34	0.096	
	5	1.84	0.24	1.5	
	6	1.10	0.38	0.038	
	7	0.32	0.53	0.032	
	8	0.36	0.38	0.0077	
	9	0.73	0.36	0.10	
	10	3.28	0.39	2.8	
	11	1.45	0.43	0.028	
	12	1.43	0.53	0.17	
	13	3.18	0.34	2.3	
	14	0.84	0.60	0.26	
	15	1.15	0.32	1.2	

 The percentage of DM particles in the Solar neighborhood originating from the LMC is small.

Local dark matter speed distribution

In the galactic rest frame



The LMC impacts the high speed tail of the DM speed distribution not only at its pericenter approach and the present day, but also up to ~175 Myr after the present day.

Direct detection event rate

• The differential event rate (per unit detector mass):



• For standard spin-independent and spin-dependent interactions:

$$\frac{dR}{dE_R} = \frac{\sigma_0 F^2(E_R)}{2m_\chi \mu_{\chi N}^2} \rho_\chi \eta(v_{\min}, t)$$

Direct detection event rate

• The differential event rate (per unit detector mass):



• For standard spin-independent and spin-dependent interactions:

$$\frac{dR}{dE_R} = \frac{\sigma_0 F^2(E_R)}{2m_{\chi}\mu_{\chi N}^2} \frac{\text{astrophysics}}{\rho_{\chi} \eta(v_{\min}, t)}$$

where

$$\eta(v_{\min}, t) \equiv \int_{v > v_{\min}} d^3 v \, \frac{f_{det}(\mathbf{v}, t)}{v}$$

Halo integral

Halo integrals



Two effects: High speed LMC particles in the Solar region + Milky Way's response to the LMC.
 Shift of > 150 km/s in the high speed tail of the halo integrals at the present day.

Direct detection exclusion limits

• Simulate the signals in 3 idealized near future direct detection experiments that would search for nuclear or electron recoils.



Xenon based detector:

Fix $\rho_{\chi} = 0.3 \text{ GeV/cm}^3$





Xenon based detector:

Fix $\rho_{\chi} = 0.3 \text{ GeV/cm}^3$





Xenon based detector:

Fix $\rho_{\chi} = 0.3 \text{ GeV/cm}^3$





Germanium based detector:





Smith-Orlik et al., JCAP 10, 070 (2023)

Direct detection: electron recoils

Silicon CCD detector:

Fix $\rho_{\chi} = 0.3 \text{ GeV/cm}^3$



Smith-Orlik et al., JCAP 10, 070 (2023)



• The LMC's influence on the local DM velocity distribution is significant even in a fully cosmological simulation.



- The LMC's influence on the local DM velocity distribution is significant even in a fully cosmological simulation.
- Our particular Sun-LMC geometry maximizes LMC's impact.

Summary

- The LMC's influence on the local DM velocity distribution is significant even in a fully cosmological simulation.
- Our particular Sun-LMC geometry maximizes LMC's impact.
- Two effects cause a boost in the high speed tail of the DM velocity distribution:
 - High speed DM particles from the LMC in the Solar region
 - The response of the Milky Way DM particles to the LMC

Summary

- The LMC's influence on the local DM velocity distribution is significant even in a fully cosmological simulation.
- Our particular Sun-LMC geometry maximizes LMC's impact.
- Two effects cause a boost in the high speed tail of the DM velocity distribution:
 - High speed DM particles from the LMC in the Solar region
 - The response of the Milky Way DM particles to the LMC

Significant shifts in direct detection limits towards lower cross sections and smaller DM masses.

Backup Slides

Local dark matter speed distribution

In the galactic rest frame (present day)



- The speed distribution of DM particles originating from the LMC peaks at the high speed tail of the Milky Way's DM distribution.
- Large halo-to-halo scatter in the results.

Changes in the halo integrals

Quantify the changes in the tails of the halo integrals by:



Impact of the DM particles from the LMC

Correlations between the percentage of LMC DM particles in the Solar region (κ_{LMC}) and $\Delta \eta$:



- $\Delta \eta$ for best fit Sun's position close to max $\Delta \eta$, and increases with κ_{LMC} .
- Scatter in $\Delta \eta$ for halos with similar $\kappa_{\rm LMC}$, due to the choice of the Sun's position for specifying the Solar region.

Impact of the DM particles from the LMC

 $\Delta \eta$ for best fit Sun's position for different snapshots in one halo:



Smith-Orlik et al., 2302.04281